

Notes 5.4 – Particle Motion Problems

Particle motion problems deal with particles that are moving along the x – or y – axis. Thus, we are speaking of horizontal or vertical movement. The position, velocity, or acceleration of a particle's motion are DEFINED by functions, but the particle DOES NOT move along the graph of the function. It moves along an axis. Most of the time, we speak of movement along the x – axis. In units 6 and 7, particle motion is revisited. At that time, we will deal more with vertical motion. For the time, we will focus on horizontal motion of particles.

In this lesson, we develop the ideas of velocity and acceleration in terms of position. We will speak of two types of velocities and accelerations. Let's define average and instantaneous velocity in the box below.

Average and Instantaneous Velocity

A particle's position is given by the function $p(t) = e^t \sin t$, where $p(t)$ is measured in centimeters and t is measured in seconds. Answer the following questions.

What is the average velocity on the interval $t = 1$ to $t = 3$ seconds? Indicate appropriate units of measure.

What is the instantaneous velocity of the particle at time $t = 1.5$. Indicate appropriate units of measure.

Before we proceed, a connection needs to be made. When given a function, $f(x)$, how did we find the slope of the secant line on the interval from $x = a$ to $x = b$? In terms of position of a particle, to what does the slope of the secant line correspond? To what does the instantaneous velocity correspond?

Average and Instantaneous Acceleration

A particle's position is given by the function $p(t) = e^t \sin t$, where $p(t)$ is measured in centimeters and t is measured in seconds. Answer the following questions.

What is the average acceleration on the interval $t = 1$ to $t = 3$ seconds? Indicate appropriate units of measure.

What is the instantaneous acceleration of the particle at time $t = 1.5$.

In summary, let's correlate the concepts of position, velocity, and acceleration to what we already know about a function and its first and second derivative.


<input type="text"/>	corresponds with	<input type="text"/>
<input type="text"/>	corresponds with	<input type="text"/>
<input type="text"/>	corresponds with	<input type="text"/>

Let's summarize our relationships between position, velocity and acceleration below.

<i>Velocity</i>	<i>Position (Motion of the Particle)</i>
Is = 0 or is undefined	
Is > 0	
Is < 0	
Changes from positive to negative	
Changes from negative to positive	

<i>Acceleration</i>	<i>Velocity</i>
Is = 0 or is undefined	
Is > 0	
Is < 0	
Changes from positive to negative	
Changes from negative to positive	

The graph below represents the position, $s(t)$, of a particle which is moving along the x axis.



- At which point(s) is the velocity equal to zero? Justify your answer.
- At which point(s) does the acceleration equal zero? Justify your answer.
- On what interval(s) is the particle's velocity positive? Justify your answer.
- On what interval(s) is the particle's velocity negative? Justify your answer.
- On what interval(s) is the particle's acceleration positive? Justify your answer.
- On what interval(s) is the particle's acceleration negative? Justify your answer.

Five Commandments of Particle Motion

1.

2.

3.

4.

5.


Suppose the velocity of a particle is given by the function $v(t) = (t + 2)(t + 4)^2$ for $t \geq 0$, where t is measured in minutes and $v(t)$ is measured in inches per minute. Answer the questions that follow.

- a. Find the values of $v(3)$ and $v'(3)$. Based on these values, describe the speed of the particle at $t = 3$.

- b. On what interval(s) is the particle moving to the left? Right? Show your analysis and justify your answer.

1998 AP Calculus AB #3 (Modified)

The graph of the velocity $v(t)$, in feet per second, of a car traveling on a straight road, for $0 \leq t \leq 50$ is shown below. A table of values for $v(t)$, at 5 second intervals of time, is also shown to the right of the graph.



t (seconds)	$v(t)$ (feet per second)
0	0
5	12
10	20
15	30
20	55
25	70
30	78
35	81
40	75
45	60
50	72

a.

During what interval(s) of time is the acceleration of the car positive? Give a reason for your answer.


b. Find the average acceleration of the car over the interval $0 \leq t \leq 50$. Indicate units of measure.

c. Find one approximation for the acceleration of the car at $t = 40$. Show the computations you used to arrive at your answer. Indicate units of measure.

d. Is the speed of the car increasing or decreasing at $t = 40$? Give a reason for your answer.

2000 AP Calculus AB #2 (Partial)

Two runners, A and B , run on a straight racetrack for $0 \leq t \leq 10$ seconds. The graph below, which consists of two line segments, shows the velocity, in meters per second, of Runner A . The velocity, in meters per second, of Runner B is given by the function v defined by $v(t) = \frac{24t}{2t+3}$.



- Find the velocity of Runner A and the velocity of Runner B at $t = 2$ seconds. Indicate units of measure.
- Find the acceleration of Runner A and the acceleration of Runner B at time $t = 2$ seconds. Indicate units of measure.

2002 AP Calculus AB #3 (Partial)

An object moves along the x – axis with initial position $x(0) = 2$. The velocity of the object at time $t \geq 0$ is given by the function $v(t) = \sin\left(\frac{\pi}{3}t\right)$.

a. What is the acceleration of the object at time $t = 4$?

b. Consider the following two statements.

Statement I: For $3 < t < 4.5$, the velocity of the object is decreasing.


Statement II: For $3 < t < 4.5$, the speed of the object is decreasing.

Are either or both of these statements correct? For each statement, provide a reason why it is correct or not correct.

Homework 5.4

A particle moves along the x axis such that its position, for $t > 0$, is given by the function $p(t) = e^{2t} - 5t$. Use this information to complete exercises 1 – 4.

1. What are the values of $p'(2)$ and $p''(2)$? Explain what each value represents.
2. Based on the values found in part (a), what can be concluded about the speed of the particle at $t = 2$? Give a reason for your answer.
3. On what interval(s) of t is the particle moving to the left? To the right? Justify your answers.
4. Does the particle ever change directions? Justify your answer.
5. The graph of $v(t)$, the velocity of a moving particle, is given below. What conclusions can be made about the movement of the particle along the x – axis and the acceleration, $a(t)$, of the particle for $t > 0$? Give reasons for your answers.



6. If the position of a particle is defined by the function $x(t) = t^3 - 9t^2 + 24t$ for $t > 0$, is the speed of the particle increasing or decreasing when $t = 2.5$? Justify your answer.

The position of a particle is given by the function $p(t) = (2t - 3)e^{2-t}$ for $t > 0$. Answer questions 7 – 9.

7. What is the average velocity from $t = 1$ to $t = 3$?
8. Find an equation for $v(t)$, the velocity of the particle.
9. For what value(s) of t will the $v(t) = 0$?

2003 AP Calculus AB #2 (Partial)

A particle moves along the x – axis so that its velocity at time t is given by

$$v(t) = -(t + 1) \sin\left(\frac{t^2}{2}\right).$$

10. Find the acceleration of the particle at $t = 2$. Is the speed of the particle increasing at $t = 2$? Explain why or why not.
11. Find all times in the open interval $0 < t < 3$ when the particle changes direction. Justify your answer.

